



A Theorem on Holonomy

Author(s): W. Ambrose and I. M. Singer

Source: *Transactions of the American Mathematical Society*, Vol. 75, No. 3, (Nov., 1953), pp. 428-443

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/1990721>

Accessed: 23/05/2008 08:59

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We enable the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.

A THEOREM ON HOLONOMY

BY

W. AMBROSE AND I. M. SINGER

Introduction. The object of this paper is to prove Theorem 2 of §2, which shows, for any connexion, how the curvature form generates the holonomy group. We believe this is an extension of a theorem stated without proof by E. Cartan [2, p. 4]. This theorem was proved after we had been informed of an unpublished related theorem of Chevalley and Koszul. We are indebted to S. S. Chern for many discussions of matters considered here.

In §1 we give an exposition of some needed facts about connexions; this exposition is derived largely from an exposition of Chern [5] and partly from expositions of H. Cartan [3] and Ehresmann [7]. We believe this exposition does however contain one new element, namely Lemma 1 of §1 and its use in passing from H. Cartan's definition of a connexion (the definition given in §1) to E. Cartan's structural equation.

1. Basic concepts. We begin with some notions and terminology to be used throughout this paper. The term "differentiable" will always mean what is usually called "of class C^∞ ." We follow Chevalley [6] in general for the definition of tangent vector, differential, etc. but with the obvious changes needed for the differentiable (rather than analytic) case. However if ϕ is a differentiable mapping we use ϕ again instead of $d\phi$ and $\delta\phi$ for the induced mappings of tangent vectors and differentials. If M is a manifold, by which we shall always mean a differentiable manifold but which is not assumed connected, and $m \in M$, we denote the tangent space to M at m by M_m . If W is a vector field at M we denote its value at m by $W(m)$, and if x_1, \dots, x_k is a coordinate system of M we always write X^1, \dots, X^k for the corresponding partial derivatives, i.e. $X^i = \partial/\partial x_i$. We use the word "diffeo" for a 1:1 differentiable mapping of one manifold onto another whose inverse is also differentiable and call the manifolds diffeomorphic. If W, W' are vector fields we write $[W, W']$ for $WW' - W'W$ (opposite to [6]). R^k will always denote k -dimensional Euclidean space of all k -tuples of real numbers and $\delta_1, \dots, \delta_k$ will always denote the canonical unit elements there, i.e. $\delta_j = (\delta_{1j}, \delta_{2j}, \dots, \delta_{nj})$.

Let G be a Lie group (i.e. a differentiable group) acting differentiably and effectively on a manifold F . If $f \in F$ and $g \in G$ we write gf for the image of f under the action of $g \in G$ and if t is a tangent vector at f we write gt for the image of t (which will be a tangent vector of gf) under g . If O is a manifold we denote by (O, F, G) the family of all transformations of $O \times F \rightarrow O \times F$ of the form: $(o, f) \rightarrow (o, t(o, f))$ when t is any differentiable mapping of $O \times F \rightarrow F$

Received by the editors September 22, 1952.

such that, for each $o \in O$, the transformation: $f \rightarrow t(o, f)$ coincides with the action of some $g \in G$.

DEFINITION. A *bundle* is a collection (B, M, F, G, π, ϕ) where B, M, F are manifolds, G is a Lie group acting differentiably and effectively on F , π is a differentiable mapping of B onto M , and ϕ is a collection of mappings. It is further assumed that

- (1) each $\phi \in \Phi$ is a diffeo of some $O \times F$, where O is an open submanifold of M , onto $\pi^{-1}(O)$,
- (2) $(\pi \circ \phi)(o, f) = o$ for all $(o, f) \in O \times F$,
- (3) if $\phi, \psi \in \Phi$, $\phi: O \times F \rightarrow B$, $\psi: P \times F \rightarrow B$, $0 \# P^{(1)}$, then $(\phi^{-1} \circ \psi) | ((O \cap P) \times F) \in (O \cap P, F, G)$,
- (4) the domains of the ϕ in Φ cover B ,
- (5) the family Φ is maximal with respect to (1), (2), (3).

Then B is called the *bundle space*, M the *base space*, G the *group*, π the *projection*, Φ the family of *strip maps* of the bundle. In the special case where F and G are the same manifold and G acts as itself by left translation, the bundle is called a *principal bundle* and denoted by (B, M, G, π, Φ) .

If ϕ is a strip map taking $O \times F \rightarrow B$, then for each $m \in O$ we define $\phi_m: F \rightarrow B$, by $\phi_m(f) = \phi(m, f)$. If ϕ, ψ are strip maps and $m \in M$ is such that both ϕ_m and ψ_m are defined, then the definition of (O, F, G) and (3) show $\phi_m^{-1} \circ \psi_m$ coincides with the action of some $g \in G$ on F and in the case of a principal bundle this means that $\phi_m^{-1} \circ \psi_m$ is a left translation by some $g \in G$.

DEFINITION. A tangent vector t at a point in a bundle is *vertical* $\text{ff}^{(2)}$ $\pi t = 0$. The linear space of vertical vectors at $b \in B$ will be denoted by V_b .

A vector field A in the fiber F which is invariant under the action of G gives rise, in a way we shall now describe, to a vertical vector field Q on B (by a vertical vector field we mean one that assigns a vertical vector at each point). To define $Q(b)$ consider any strip map $\phi: O \times F \rightarrow B$, such that $b \in \phi(O \times F)$. If $\phi_m(f) = b$ we define $Q(b) = \phi_m A(f)$. This definition is independent of the strip map ϕ because if ψ is another such strip map then $\phi_m^{-1} \circ \psi_m$ agrees with the action of some $g \in G$. Hence $\psi_m(f) = \phi_m(gf)$ for all $f \in F$ and then, using the invariance of A , we have $\psi_m A(f) = \phi_m g A(f) = \phi_m A(gf)$. If A is differentiable, then so is W and the mapping thus defined, taking $A \rightarrow W$, preserves brackets (these properties are immediate from the fact that the strip maps ϕ are diffeos). In particular, if the bundle is a principal bundle, the invariant vector fields on the fiber are the left invariant vector fields of the group G and these constitute the Lie algebra L of G . Hence in this case the mapping taking $A \rightarrow W$ is an iso⁽³⁾ of the Lie algebra L of G onto a Lie algebra of vertical vector fields on B . Throughout this paper we shall,

(1) $0 \# P$ means 0 and P have a non-null intersection. Similarly $0 || P$ will mean 0 and P have a null intersection.

(2) ff means: if and only if.

(3) We shall omit the appendage "morphism" from an obvious maximal set of words.

in the case of a principal bundle (and principal bundles will be the only ones we shall seriously consider), reserve the letter q for this iso and the letter \mathcal{Q} for its image $q(L)$. If t is a vertical vector at $b \in B$, then it is trivial that there exists a unique $Q \in \mathcal{Q}$ such that $Q(b) = t$; we then say Q is *generated* by t and if $Q = q(A)$ we also say A is *generated* by t .

If (B, M, G, π, Φ) is a principal bundle, then G can be made to act on B in a natural way that is called *right translation*. This differentiable and effective action of G on B is defined as follows. If $b \in B, g \in G, \phi$ any strip map, $\phi: O \times F \rightarrow B, b \in \phi(O \times F)$, then write $b = \phi(m, h)$ and define $bg = \phi(m, hg)$. This definition of bg is independent of the choice of the strip map ϕ because, for any other strip map $\psi, \phi_m^{-1} \circ \psi_m$ is a left translation on G and left translations commute with right translations on G . The right translation action of G on B is the mapping of $B \times G \rightarrow B: (b, g) \rightarrow bg$. The mapping of $B \rightarrow B$ taking $b \rightarrow bg$ (for any fixed $g \in G$) will be denoted by R_g and right translation by g on G will also be denoted by R_g ; left translation by g on G will be denoted by L_g . If ϕ is any strip map of B , then clearly $\phi_m \circ R_g = R_g \circ \phi_m$.

We denote the inner auto⁽³⁾ of G , corresponding to $g \in G$, by $\alpha(g)$, i.e. $\alpha(g)h = ghg^{-1}$ for $h \in G$, and if $A \in L$ (L the Lie algebra of G) we write $\alpha(g)A$ for the element of L defined by: $(\alpha(g)A)(e) = \alpha(g)A(e)$. $\alpha(g)A$ is sometimes denoted by $adg A$. For $Q \in \mathcal{Q}$ we define $\alpha(g)Q \in \mathcal{Q}$ by $\alpha(g)Q = q(\alpha(g)A)$, when $Q = q(A)$. Then $R_gQ = \alpha(g^{-1})Q$, for if ϕ is any strip map of B and $\phi(h) = b$ then $(\alpha(g^{-1})Q)(b) = \phi((\alpha(g^{-1})A)(h)) = \phi(L_h \circ R_g \circ L_{g^{-1}}A(e)) = \phi(R_g \circ L_{hg^{-1}}A(e)) = \phi(R_gA(hg^{-1})) = (\phi \circ R_g)A_{(hg^{-1})} = (R_g \circ \phi)(A(hg^{-1})) = R_gQ(hg^{-1})$.

Some principal bundles of special interest are the *bundle of bases* $E(M)$ over any manifold M and the *bundle of frames* $F(M)$ over a Riemannian M . We now define $E(M)$. The points of $E(M)$ are all the ordered $(d+1)$ -tuples (m, e_1, \dots, e_d) where $m \in M$ and (e_1, \dots, e_d) is any base of M_m . The projection π of $E(M)$ onto M is defined by $\pi(m, e_1, \dots, e_d) = m$. The group (and fiber) G is the full linear group over the reals, i.e. the group of all nonsingular linear transformations of R^d onto itself. With each point b of $E(M)$ we can associate an iso I_b of M_m onto R^d by writing $b = (m, e_1, \dots, e_d)$ and defining I_b to be the linear transformation taking $e_i \rightarrow \delta_i$ (for all i); and conversely, so the points of $E(M)$ could have been defined as the pairs (m, I) where I is such an iso. In this example it is convenient to define the action of G on the bundle before defining the bundle structure. We do this by: if $b = (m, e_1, \dots, e_d)$, then $bg = (m, I_b^{-1}gI_b e_1, \dots, I_b^{-1}gI_b e_d)$; this implies that I_{bg} takes $(I_b^{-1} \circ g \circ I_b)e_i \rightarrow f_i$, hence $I_{bg} \circ I_b^{-1} \circ g \circ I_b = I_b$ from which it is clear that $I_{bg} = g^{-1} \circ I_b$ (a fact that will be useful later). For any coordinate system x_1, \dots, x_d of M with domain O we define $\phi: O \times G \rightarrow E(M): (m, g) \rightarrow R_g(m, X'(m), \dots, X^d(m))$ ⁽⁴⁾ (now writing $R_g b$ for the bg just defined). We introduce coordinates $y_1, \dots, y_d, y_{d+1}, \dots, y_{dd}$ of $E(M)$, with domain $\pi^{-1}(O)$, by

⁽⁴⁾ The notation $f: X \rightarrow Y: x \rightarrow y$ means f is a function from X to Y taking x into y .

$$y_j = x_j \circ \phi, \quad e_j = \sum_i y_{ij}(m, e_1, \dots, e_d) X^i(m).$$

Such coordinate systems of $E(M)$ obtained from different coordinate systems of M are easily seen to be differentiably related. Since their domains cover $E(M)$ they define a unique differentiable structure on $E(M)$, and this defines $E(M)$ as a manifold. Now let x_1, \dots, x_d and x'_1, \dots, x'_d be any two coordinate systems of M with domains O and O' , let ϕ and ϕ' be the mappings they define as above, and we shall show $(\phi^{-1} \circ \phi')|((O \cap O') \times G)$ is in $(O \cap O', G, G)$. If $m \in O \cap O'$, $b = \phi(m, e) = (m, X^1(m), \dots, X^d(m))$, $b' = \phi'(m, e) = (m, X'^1(m), \dots, X'^d(m))$, then $\phi_m^{-1} \phi'_m$ takes $g \in G$ into the $h \in G$ such that $I_b^{-1} g I_b X^i(m) = I_{b'}^{-1} h I_{b'} X'^i(m)$, i.e. g goes into the h such that $I_b^{-1} g \delta_i = I_{b'}^{-1} h \delta_i$ (for all i). Hence $h = I_{b'} \circ I_b^{-1} \circ g$, showing that $\phi_m^{-1} \circ \phi'_m$ is left translation by $k = I_{b'} \circ I_b^{-1}$ on G . And from this expression for k it is easily seen that k is a differentiable function of m , thus $(\phi^{-1} \circ \phi')|((O \cap O') \times G)$ is in $(O \cap O', G, G)$. We define Φ , the family of strip maps of this bundle, to be all ϕ obtained from coordinate systems of M as above plus all other mappings ψ of any $P \times G \rightarrow E(M)$ which satisfy conditions (1) and (2) in the definition of a bundle and the condition that $(\psi^{-1} \circ \phi)|((O \cap P) \times G)$ is in $(O \cap P, G, G)$ for such ϕ as above. Clearly the R_g defined by this bundle structure agrees with the R_g used above.

The bundle $F(M)$ is defined similarly, the points in this case being of the form (m, e_1, \dots, e_d) where $m \in M$ and the e_i are any orthonormal base of M_m . $F(M)$ is in a natural way a submanifold of $E(M)$.

DEFINITION. A connexion H on a principal bundle $\mathcal{B} = (B, M, G, \pi, \Phi)$ is a mapping which assigns to each $b \in B$ a linear subspace H_b of B_b such that

- (1) H_b is a linear complement to V_b ; elements of H_b will be called *horizontal* vectors.
- (2) H is invariant under the action of G on B , i.e. $H_{bg} = R_g H_b$,
- (3) H_b depends differentiably on b , i.e. if W is any differentiable vector field on B , HW its horizontal part (i.e. HW is the vector field which assigns at each $b \in B$ the projection of $W(b)$ into H_b), HW is differentiable.

NOTATION. If W is any vector field on such a B with a connexion H we shall always write HW and VW for the horizontal and vertical parts of W ; differentiability of W trivially implies that of VW .

If H is a connexion on B then, for each $b \in B$, $\pi|H_b$ is an iso of H_b onto M_m , where $m = \pi b$. This is immediate from: (1) π maps B_b onto M_m , (2) $\pi V_b = 0$, (3) $\dim H_b = \dim M_m$.

DEFINITION. If H is a connexion on B and $b \in B$, the *holonomy group* of H attached to b , denoted by $G(b)$, is the set of all $g \in G$ such that bg can be joined to b by a piecewise-differentiable⁽⁵⁾ horizontal curve in B . The *null holonomy*

(5) By a *piecewise differentiable* curve we mean a curve which is differentiable and has a nonzero tangent vector except at a finite number of points. By *horizontal* we mean that the tangent vector is in H_b at all points b on the curve for which the tangent vector exists.

group of H attached to b , denoted by $G_0(b)$, is the set of all $g \in G$ such that bg can be joined to b by a piecewise differentiable horizontal curve in B whose projection into M is null-homotopic.

It is trivial that $G(b)$ and $G_0(b)$ are subgroups of G and that if $c = bg$ then $G(c) = g^{-1}G(b)g$, $G_0(c) = g^{-1}G_0(b)g$.

If H is a connexion on $E(M)$, then the holonomy group is related to the notion of parallel translation along curves in M so we briefly indicate here how parallel translation is derived from the above concept of a connexion. Let H be a connexion on $E(M)$. If ρ is any piece-wise differentiable curve in M , from m to m' , we shall define a linear transformation T of M_m onto $M_{m'}$ and this linear transformation is called *parallel translation along ρ* . To define T let σ be any horizontal curve in $E(M)$ lying over ρ (i.e. $\rho = \pi \circ \sigma$). Then σ starts at some point (m, e_1, \dots, e_d) and ends at some point (m', e'_1, \dots, e'_d) . We define T to be the unique linear transformation taking $e_i \rightarrow e'_i$ (for all i). The horizontal curve σ lying over ρ is not unique, but because of the invariance of H under the action of G on $E(M)$ it is easily seen that T is independent of the choice of σ . The existence of a horizontal curve σ lying over ρ can be shown as follows (note that this construction is valid for any connexion, not necessarily on an $E(M)$). Consider any curve τ lying over ρ in B —and such curves trivially exist by virtue of the local cross section given by strip maps. Consider the submanifold B' of B consisting of all fibers which contain a point of τ , with the natural differentiable structure in which it is locally the product space of the fiber by the parameter along τ . We define a vector field W on B' by translating the tangent vector to τ up and down each fiber in B' under the action of G on B . Then HW is a vector field on B' any of whose integral curves will be such a σ . In terms of parallel translation we thus can consider the holonomy group of a connexion on $E(M)$ to be a group of linear transformations on the tangent space at a point of M .

A connexion H on B gives rise to a 1-form ω , the values of ω lying in the Lie algebra L of the group G of the bundle. ω is defined by: if $t \in B_b$ then $\omega(t) =$ the element of L generated by Vt , i.e. $\omega(t)$ is the unique $A \in L$ with $(qA)_b = Vt$. Clearly ω has the following properties.

- (1) ω is differentiable⁽⁶⁾,
- (2) if t is vertical, then $\omega(t)$ is the element of L generated by t ,
- (3) $\omega \circ R_g = (adg^{-1})\omega$, i.e. if $t \in B_b$ then $\omega(R_g t) = (adg^{-1})(\omega(t))$.

We have (3) because $VR_g t = R_g Vt$ and $R_g A = (adg^{-1})A$ for $A \in L$. Conversely, any 1-form ω on B with values in L and satisfying (1), (2), (3) gives rise to a connexion H defined by: $H_b = [t \in B_b | \omega(t) = 0]$. We call ω the *1-form of the connexion*.

In differentiating a real-valued differentiable differential p -form θ we shall frequently make use of the formula:

⁽⁶⁾ For a form ω whose values lie in a finite-dimensional linear space over R , differentiability means that for every linear functional f from the vector space to R , $f \circ \omega$ is differentiable.

$$\begin{aligned}
 d\theta(W_1, \dots, W_{p+1}) &= \frac{1}{p+1} \sum_{i=1}^{p+1} (-1)^{i-1} W_i(\theta(W_1, \dots, \widehat{W}_i, \dots, W_{p+1})) \\
 &\quad + \frac{1}{p+1} \sum_{i < j} (-1)^{i+j} ([W_i, W_j], W_1, \dots, \widehat{W}_i, \dots, \widehat{W}_j, \dots, W_{p+1})
 \end{aligned}$$

which can be used to define $d\theta$. I.e. this formula holds for any $p+1$ differentiable vector fields W_1, \dots, W_{p+1} , and if this formula holds at a point for all choices of W_1, \dots, W_{p+1} from a family of vector fields which span the tangent space at that point, then this determines $d\theta$ at the point. If θ is a differentiable differential form with values in a finite-dimensional vector space over R , we define $d\theta$ in the usual way via linear functionals, i.e. $d\theta$ is the unique form such that $f \circ d\theta = d(f \circ \theta)$ for all such linear functionals f .

DEFINITION. If H is a connexion or a principle bundle with bundle space B we define the *covariant derivative* $D\theta$ of a differentiable p -form θ on B (when the values of θ lie in any finite-dimensional vector space over R) to be the $(p+1)$ -form on B defined by

$$(D\theta)(t_1, \dots, t_{p+1}) = d\theta(Ht_1, \dots, Ht_{p+1})$$

where t_1, \dots, t_p are any elements of B_b . I.e. D is the exterior derivative followed by the dual of the projection H . Clearly $D\theta$ is differentiable.

DEFINITION. If H is a connexion whose 1-form is ω , we define the curvature form Ω of the connexion by $\Omega = D\omega$.

It is easily seen that $\Omega \circ R_g = (adg^{-1})\Omega$, i.e. $\Omega(R_g t_1, R_g t_2) = adg^{-1}\Omega(t_1, t_2)$ for any $t_1, t_2 \in B_b$. This follows from the corresponding fact for ω and the facts (which are verified by simple computations) that if a form θ with values in L satisfies this relation, then so do $d\theta$ and $H^*\theta$, where H^* is the dual of H , i.e. $(H^*\theta)(t_1, \dots, t_p) = \theta(Ht_1, \dots, Ht_p)$.

We now explain, in the case of a bundle of bases $E(M)$, the geometric reason for calling this Ω the curvature. We show that Ω defines, at each $m \in M$ and for each $s, s' \in M_m$, a linear transformation $T_{s,s'}$, where this transformation can be considered as the effect of translating vectors parallel to themselves around the "infinitesimal parallelogram" spanned by s and s' ; so this transformation, as a function of s and s' , is a measure of how M "curves" at m —this "curving" depending on the connexion. We also point out in the case of a Riemannian connexion how to get the Riemannian curvature, which is a number depending on m, s, s' , from $T_{s,s'}$.

Let $s, s' \in M$, consider any b in $E(M)$ lying over m (i.e. $\pi b = m$) and any horizontal differentiable vector fields K and K' defined on a neighborhood of b such that $\pi K(b) = s, \pi K'(b) = s'$. We define for each small positive number δ a "circuit" γ_δ , which is a piecewise differentiable horizontal curve consisting of four differentiable pieces, as follows. Take the integral curve ρ of K with

$\rho(0) = b$, then an integral curve ρ' of K' with $\rho'(0) = \rho(\delta)$, then an integral curve σ of K with $\sigma(0) = \rho'(\delta)$, then an integral curve σ' of K' with $\sigma'(0) = \sigma(-\delta)$. The circuit γ_δ will be the curve obtained by going along ρ from $\rho(0)$ to $\rho(\delta)$, then along ρ' from $\rho(\delta) = \rho'(0)$ to $\rho'(\delta)$, then along σ from $\rho'(\delta) = \sigma(0)$ to $\sigma(-\delta)$, then along σ' from $\sigma(-\delta) = \sigma'(0)$ to $\sigma'(-\delta)$. Next we consider the curve made up of the end points of these circuits, suitably parametrized, i.e. we define $\gamma(t) = \text{final point of } \gamma_{t^{1/2}}$. (This is our definition of γ for $t > 0$; for $t < 0$ we can define it similarly or we can ignore all $t < 0$.) If we define $\gamma(0) = b$, then γ is a differentiable curve whose tangent vector at b is $[K, K'](b)$ (this is just the geometrical significance of the bracket operation, applied to our K and K'). Let η_δ and η be the projection of γ_δ and γ into M . Note that η_δ is a closed curve if and only if the final point of γ_δ lies on the same fiber as b and in this case $b = (m, e_1, \dots, e_d)$, the final point of γ_δ is (m, f_1, \dots, f_d) , and parallel translation around η_δ is the linear transformation taking $e_i \rightarrow f_i$; this parallel translation gives rise to an element $g \in G$ (G the linear group), namely the g such that $R_g(m, e_1, \dots, e_d) = (m, f_1, \dots, f_d)$. In general η_δ is not closed and the final point of γ_δ does not lie on the same fiber as b , however the vertical component of the tangent vector to γ at b is an infinitesimal analogue of the transformation mentioned above carrying $e_i \rightarrow f_i$. The element A of L (L the Lie algebra of G) generated by the vertical component of the tangent vector to γ at b is an infinitesimal analogue of the above g and we point out in the next paragraph how, through A , the vertical component of this tangent gives rise to a transformation $T_{s,s'}$ on M_m . This $T_{s,s'}$, as a function of s, s', m (s and s' in M_m), determines the curvature form Ω and conversely.

By definition of the 1-form ω of the connexion we have $A = \omega([K, K'](b))$ and a simple calculation (which is given in the proof of Theorem 2 below) shows $\omega([K, K'](b)) = -\Omega(K(b), K'(b))$. Thus we have shown that if t and t' are the unique horizontal elements of B_b which project to s and s' (we can discard the vector fields K and K' at this point), then the element A of L generated by the vertical component of the tangent vector to γ at b is given by $A = -\Omega(t, t')$. Because G is the linear group, each $A \in L$ gives rise to a linear transformation T_A on R^d defined by: $f(T_A x) = A(e)f_x$, where f is any linear function on R^d and f_x is the function defined on G by $f_x(g) = f(gx)$, for $x \in R^d$. (In terms of the usual coordinates x_{ij} on G this just defines T_A as the linear transformation whose matrix with respect to the base $\delta_1, \dots, \delta_d$ is $(A(e)x_{ij})$.) We define $T_{s,s'} = I_b^{-1} \circ T_A \circ I_b$, where I_b is the mapping defined above of M_m onto R^d , associated with b . It remains only to show this definition is independent of the point b over m —a consequence of the behavior of Ω under R_g . To prove this let bg be another point over m , t_1 and t'_1 the horizontal vectors at bg that project to s and s' , so $R_g t = t_1$, $R_g t' = t'_1$, and let $A_1 = -\Omega(t_1, t'_1)$. Because $\Omega(t_1, t'_1) = \Omega(R_g t, R_g t') = \text{ad}g^{-1}\Omega(t, t')$ we have

$T_A = g^{-1}T_Ag$. Then using that $I_{bg} = g^{-1} \circ I_b$ we have $I_{bg}^{-1} \circ T_A \circ I_{bg} = I_b^{-1} \circ g \circ g^{-1} \circ T_A \circ g \circ g^{-1} \circ I_b = I_b^{-1} \circ T_A \circ I_b$.

Consider now a Riemannian connexion on $E(M)$ (the Riemannian connexion is usually considered on the bundle of frames $F(M)$ but may equally well be considered on $E(M)$ —or any principal bundle containing $F(M)$ as a sub-bundle). I.e. we have a scalar product \langle , \rangle on each M_m which is a differentiable function of m and H is the unique connexion on $E(M)$ with torsion zero (torsion is discussed at the end of this section) and such that parallel translation preserves the scalar product. Then the function considered above which assigns to each s and s' in M_m the linear transformation $T_{s,s'}$ can be characterized by the simpler function K which assigns to each such s and s' the number $K = \langle T_{s,s'}s, s' \rangle / \alpha(s, s')^2$ where $\alpha(s, s')$ is the “area” of the parallelogram spanned by s and s' , i.e. $\alpha(s, s') = (\|s\|^2\|s'\|^2 - \langle s, s' \rangle^2)^{1/2}$. It is easily seen that $K(s, s')$ does not depend on s and s' but only on the plane section (i.e. linear subspace of M_m) spanned by s and s' . This K is the Riemannian curvature.

For the proof of the structural equation of a connexion we need the following lemma, which might be called the vector field formulation of the structural equation.

LEMMA. *Let H be any connexion on a principal bundle with bundle space B , and \mathcal{Q} as above. If K is any horizontal vector field on B and $Q \in \mathcal{Q}$ then $[K, Q]$ is horizontal.*

Proof. We need first the fact: if A and B are vector fields on a Lie group G with A left invariant and B right invariant, then $[A, B] = 0$. This can be proved as follows. Define vector fields A' and B' on $G \times G$ by: $A'(g, h) = J_g A(h)$, $B'(g, h) = I_h B(g)$ where

$$I_h : G \rightarrow G \times G : g \rightarrow (g, h); \quad J_g : G \rightarrow G \times G; \quad h \rightarrow (g, h).$$

Then $[A', B'] = 0$ for if we consider any $(g, h) \in G \times G$ and choose coordinates x_1, \dots, x_n on a neighborhood of g , coordinates y_1, \dots, y_n on a neighborhood of h , then $A' = \sum a_i Y^i$, $B' = \sum b_i X^i$ where the a_i depend only on the y_j -coordinates and the b_i depend only on the x_j -coordinates. Computation shows $\Phi A' = A$ and $\Phi B' = B$ where Φ denotes group multiplication in G . Hence $[A, B] = 0$.

It follows that for any vertical vector field W on $\pi^{-1}(0)$ (0 any open subset of M) which is invariant under the action of G on B we have $[W, Q] = 0$ for all $Q \in \mathcal{Q}$.

Now consider any strip map $\phi : 0 \times G \rightarrow B$ where there exists a coordinate system x_1, \dots, x_d of M with domain O and let $x_j = x_j \circ \pi$. We call a coordinate system of B *special* if it consists of the functions y_1, \dots, y_d plus some functions z_1, \dots, z_k which are carried over by ϕ from a coordinate

system v_1, \dots, v_k of G (i.e. $z_j = v_j \circ p \circ \phi^{-1}$ where $p: 0 \times G \rightarrow G: (m, g) \rightarrow g$); if the domain of v_1, \dots, v_k is P , then the domain of this special system is $\phi(0 \times P)$. If we have two special coordinate systems (special, that is, for the same ϕ and x_1, \dots, x_d) then at any point b common to their domains the partial derivative with respect to y_j at b defined through the two coordinate systems is easily seen to be the same tangent vector and we denote the vector field thus defined throughout $\pi^{-1}(0)$ by Y^i , i.e. $Y^i(b)$ is the partial derivative with respect to y_j , where $y_1, \dots, y_d, z_1, \dots, z_k$ is any special coordinate system whose domain includes b . Then $\pi Y^i = X^i$ and $R_g Y^i = Y^i$. Write $Y^i = K^i + W^i$ with K^i horizontal and W^i vertical. We next show $[K^i, Q] = 0$ for all $Q \in \mathcal{Q}$, by showing separately that $[Y^k, Q] = 0$ and $[W^j, Q] = 0$.

To prove $[Y^i, Q] = 0$ consider any $b \in \pi^{-1}(0)$ and a special coordinate system $y_1, \dots, y_d, z_1, \dots, z_k$ at b . The expression for Q in these coordinates involves only the z_i and with coefficients that depend only on the z_i . Hence $[Y^i, Q] = 0$. To prove $[K^i, Q] = 0$ it is sufficient, by a remark made above, to show $R_g K^i = K^i$. Because $R_g K^i$ is horizontal, $R_g W^i$ is vertical, and the horizontal vectors at any point are a linear complement to the vertical vectors, this follows from

$$K^i + W^i = Y^i = R_g Y^i = R_g K^i + R_g W^i.$$

Now consider any horizontal vector field K defined on a neighborhood of any $b \in \phi(0 \times G)$. Using the above notation, the K^i form a base for the horizontal vectors throughout $\phi(0 \times G)$ (because of the iso of H_b with M_m and the fact that the X^i are a base for M_m) so $K = \sum a_j K^j$. Then

$$[K, Q] = [\sum a_j K^j, Q] = \sum a_j [K^j, Q] - \sum (Q a_j) K^j = - \sum (Q a_j) K^j.$$

THEOREM 1. (*The structural equation of a connexion*). *If H is any connexion on a principal bundle with bundle space B , ω is the 1-form of the connexion, and Ω is the curvature form of the connexion, then*

$$d\omega = - [\omega, \omega]/2 + \Omega,$$

i.e. for any $b \in B$ and $s, t \in B_b$ we have

$$d\omega(s, t) = - [\omega(s), \omega(t)]/2 + \Omega(s, t).$$

Proof. The formula mentioned above for the differential of a real-valued differentiable differential p -form θ becomes, in case $p = 1$,

$$d\theta(W, W') = (W\theta(W') - W'\theta(W) - \theta([W, W']))/2,$$

where W and W' are any differentiable vector fields. Further, if η is a 2-form and we wish to show $d\theta = \eta$, it is sufficient to find for each $b \in B$ a family of vector fields W^1, \dots, W^r for which the $W^i(b)$ span B_b and such that $d\theta(W^i(b), W^j(b)) = \eta(W^i(b), W^j(b))$ for all i, j .

The differential forms of our theorem have values in a Lie algebra so we must prove, for any linear functional f on that Lie algebra, that

$$f(d\omega(W, W')) = -f([\omega(W), \omega(W')])/2 + f(\Omega(W, W')).$$

Letting $\theta = f \circ \omega$ this becomes

$$d\theta(W, W') = -f([\omega(W), \omega(W')])/2 + d\theta(HW, HW')$$

and using the above formula for $d\theta$ this becomes

$$\begin{aligned} W\theta(W') - W'\theta(W) - \theta([W, W']) &= -f([\omega(W), \omega(W')]) + (HW)(\theta(HW')) \\ &\quad - (HW')(\theta(HW)) - \theta([HW, HW']). \end{aligned}$$

It is sufficient to prove this at each $b \in B$, for all choices of W and W' from any family W^1, \dots, W^r which are defined on a neighborhood of b and such that $W^i(b)$ span B_b . We choose such a family consisting of any Q^1, \dots, Q^k which span \mathcal{Q} plus any horizontal K^1, \dots, K^d for which the $K^i(b)$ span H_b .

The desired formula holds whenever W is a K^i and W' a K^i , for then $\omega(W) = 0, \omega(W') = 0$, hence $f([\omega(W), \omega(W')]) = 0$; and $W = HW, W' = HW'$, so the two sides are identical. If W is a Q^i and W' is a K^i , then $\omega(W') = 0$ hence $f([\omega(W), \omega(W')]) = 0$, and $HW = 0$ so the entire right side is zero. In this case the left side is also zero, $\theta([W, W'])$ being zero because the previous lemma shows in this case that $[W, W']$ is horizontal, $\theta(W')$ being zero because W' is horizontal, and $W'\theta(W)$ being zero because $\theta(W)$ is a constant function. If W is a Q^i and W' is a Q^j , then (because g is a Lie algebra iso) $f([\omega(W), \omega(W')]) = f(\omega[W, W']) = \theta([W, W'])$ and HW and HW' are 0 so the left side reduces to $-\theta([W, W'])$. Because both $\theta(W)$ and $\theta(W')$ are constant functions the right side also reduces to $-\theta([W, W'])$. Hence the formula is proved.

In the special case of a connexion on a bundle of bases, and if the elements of the Lie algebra of the linear group are represented in the usual way by matrices, then $\omega(t)$ is represented by $(\omega_{ij}(t))$, $\Omega(s, t)$ is represented by $(\Omega_{ij}(s, t))$, and the above formula becomes

$$d\omega_{ij} = - \sum \omega_{i\alpha} \wedge \omega_{\alpha j} + \Omega_{ij}.$$

In this case, however, one gets the further formula

$$d\omega_i = - \sum \omega_{\alpha i} \wedge \omega_\alpha + \Omega_i$$

which we now explain. One defines 1-forms $\omega_1, \dots, \omega_d$ on $E(M)$ by: if $t \in E(M)_b$ and $b = (m, e_1, \dots, e_d)$, then $\omega_i(t)$ is the i th coefficient of πt when πt is expressed linearly in terms of the base e_1, \dots, e_d of M_m . Then Ω_i is defined to be the covariant derivative of ω_i and this formula is proved similarly to the other; the Ω_i are called the *torsion forms* of the connexion. Let E^1, \dots, E^d be the vector fields defined on $E(M)$ by: if $b = (m, e_1, \dots, e_d)$, then $E^i(b)$ is the unique element of H_b that projects to e_i under π . One verifies

easily that torsion zero (i.e. all $\Omega_i = 0$) is equivalent to the statement that $[E^i, E^j]$ is vertical for all i, j . In fact the role of the Ω_i is precisely to describe the horizontal components of the $[E^k, E^r]$, just as the role of the Ω_{ij} is to describe their vertical components; the significant facts about such a connexion may be described either through the $[E^i, E^j]$ or through the Ω_{ij} and Ω_k . A more elegant way of interpreting the above formula for $d\omega_i$, however, is to pass from the bundle $E(M)$ to a larger bundle whose group is the affine group. The connexion H on $E(M)$ gives rise naturally to a connexion K on the larger bundle, and with the following properties. Let Θ be the curvature form of K . Then Θ decomposes into $\Theta' + \Theta''$ through the decomposition of the Lie algebra of the affine group into the subalgebras corresponding to the linear transformations and the translations. Θ' is essentially the curvature form of H and Θ'' is essentially the torsion. The structural equation for K can be similarly decomposed into two parts, the two parts being essentially the equations for $d\omega_{ij}$ and for $d\omega_i$.

2. Curvature as generator of the holonomy group. In this section we consider a connexion H on a principal bundle (B, M, G, π, Φ) . We assume further that M is connected and that its fundamental group has at most a countable number of elements. We denote the 1-form of the connexion by ω and the curvature form by Ω . The Lie algebra of G will be denoted by L . If $b \in B$, then the holonomy group and null holonomy group associated with b will be denoted by $G(b)$ and $G_0(b)$, while the arc-component of the identity of $G(b)$ will be denoted by $G(b)^0$. $B(b)$ will denote the subset of B consisting of all points which can be joined to b by piece-wise differentiable horizontal curves. This notation will be kept fixed throughout the section.

THEOREM 2. *For any $b \in B$ let $L(b)$ be the subalgebra of L generated by all $\Omega(s, t)$ where s and t run through all pairs of tangent vectors to B at all points of $B(b)$. Then the subgroup of G generated by $L(b)$ is $G_0(b)$.*

We prove this via a sequence of lemmas. The first two are needed to prove $B(b)$ a submanifold of B , which fact is important in the proof of this theorem.

LEMMA 1. *Let ϕ be a strip map of the bundle, $\phi = O \times G \rightarrow B$, where O is an open submanifold of M . Suppose that O is the domain of a coordinate system x_1, \dots, x_d of M with center m_0 (i.e. all $x_i(m_0) = O$) and that O is an open ball with respect to these coordinates. Let $b_0 = \phi(m_0, e)$. We now define a mapping α of $O \rightarrow G$ (α will depend on the coordinates x_i) and the assertion of this lemma is that α is differentiable on some neighborhood P of m_0 . To define α consider, for each $m \in O$, the ray ρ_m (i.e. ρ_m is a ray with respect to the coordinates x_i) from m_0 to m , parametrized so that $\rho_m(0) = m_0, \rho_m(1) = m$. Lying above ρ_m we have curves σ_m and τ_m in B defined as follows. σ_m is the unique horizontal curve in B starting at b_0 and lying over ρ_m . τ_m is defined by: $\tau_m(u) = \phi(\rho_m(u), e)$. Then $\sigma_m(1) = \tau_m(1)g$ for a unique $g \in G$ and we define $\alpha(m) = g$.*

Proof. Let v_1, \dots, v_k be any coordinate system on some neighborhood U of e in G . To prove the lemma we shall show the functions $v_j \circ \alpha$ are differentiable on some neighborhood P of m_0 . The x_i and v_j give coordinates in the usual way on $\phi(O \times G)$ which are carried by ϕ to coordinates on B ; we denote these coordinates, whose domain is $\phi(O \times U)$, by y_1, \dots, y_{k+d} .

We first prove the existence of σ_m as stated in the lemma. We define σ_m to be the integral curve with $\sigma_m(0) = b_0$ of the vector field $H(\sum_{i=1}^d x_i(m) Y^i)$ —this vector field is defined on $\pi^{-1}(0)$ and is differentiable by condition (3) in the definition of a connexion, hence σ_m is a horizontal differentiable curve. We have $\pi \circ \sigma_m = \rho_m$ because $\pi H(\sum_{i=1}^d x_i(m) Y^i) = \sum x_i(m) X^i$, which implies $\pi \circ \sigma_m$ is an integral curve of $\sum x_i(m) X^i$, as is also ρ_m , hence $\pi \circ \sigma_m = \rho_m$. Uniqueness of σ_m follows from the uniqueness of integral curves, for any such σ_m is an integral curve of $H(\sum_{i=1}^d x_i(m) Y^i)$.

If $m \in O$ we shall write um for $\rho_m(u)$. Since $\phi(m, e)g = \phi(m, g)$ we have $\phi(m, \alpha(m)) = \phi(m, e)\alpha(m) = \sigma_m(1)$, and since $\sigma_m(u) = \sigma_{um}(1)$ we then have $\phi(um, \alpha(um)) = \sigma_m(u)$. Because σ_m is a differentiable curve in B we have, for each $m \in O$, the existence of $u_m > 0$ such that $\sigma_m(u) \in U$ for $u < u_m$, and we take u_m to be the largest possible such number. Then $v_j(\alpha(um)) = y_{j+d}(\phi(um, \alpha(um))) = y_{j+d}(\sigma_m(u))$ for $u < u_m$.

Because σ_m is an integral curve of $H(\sum_{i=1}^d x_i(m) Y^i)$ we have

$$(D(y_{j+d} \circ \sigma_m))(u) = \left(dy_{j+d} \left(H \left(\sum_{i=1}^d y_i Y^i \right) \right) \right) (\sigma_m(u))$$

where D denotes the usual differentiation operator on R . From the definition of a connexion it follows that the function $dy_{j+d}(H(\sum_{i=1}^d y_i Y^i))$ is a differentiable function and hence can be expressed on a neighborhood of b_0 as a differentiable function of the y_r . This shows the functions $y_r \circ \sigma_m$ are solutions of a system of ordinary differential equations with differentiable functions on the "right side." It follows that the functions f_j defined by: $f_j(u, m) = (y_j \circ \sigma_m)(u)$ for $u < u_m$ are defined and differentiable on some neighborhood of (O, m_0) .

Because $\sigma_m(u) = \sigma_{um}(1)$ it follows that $\sigma_m(1) \in U$ for m in some neighborhood P of m_0 and then, because $(v_j \circ \alpha)(m) = y_j(\sigma_m(1)) = f_j(1, m)$, we have that $v_j \circ \alpha$ is differentiable on P .

LEMMA 2. Let $m_0 \in M$, b_0 be any point of B lying over m_0 , and x_1, \dots, x_d any coordinate system of M whose domain includes m_0 and has center at m_0 . Then there exists a strip map $\psi: P \times G \rightarrow B$, where P is an open ball with respect to the coordinates x_i , such that $\psi(m_0, e) = b$ and for every ray ρ_m from m_0 to m in P (with respect to these coordinates) the curve $\psi(\rho_m, e)$ in B is horizontal. We call such a strip map canonical with respect to the coordinates x_i .

Proof. Let ϕ be any strip map taking $O \times G \rightarrow B$, with $\phi(m_0, e) = b_0$, where O is an open subset of the domain of the coordinates x_i . We define α , depending

on ϕ and the x_i , as in the previous lemma. Let P be any neighborhood of m_0 on which α is differentiable and such that P is a ball with respect to the coordinates x_i . We define ψ by: $\psi(m, g) = \phi(m, \alpha(m)g)$. ψ is differentiable because α and ϕ are differentiable. ψ is a strip map because, for any strip map χ , $(\chi_m^{-1} \circ \psi_m)(g) = (\chi_m^{-1} \circ \phi_m)(\alpha(m)g)$, so if $\chi_m^{-1} \circ \phi_m = L_h$ ⁽⁷⁾ then $\chi_m^{-1} \circ \psi_m = L_h\alpha(m)$. And $\psi(\rho_m, e)$ is horizontal because $\psi(\rho_m(u), e) = \phi(\rho_m(u), \alpha(\rho_m(u))) = R\alpha(\rho_m(u))\phi(\rho_m(u), e)$.

We now define the differentiable structure in which $B(b)$ is a submanifold of B . First note that $G(b)^0$, being an arcwise connected subgroup of the Lie group G , is a differentiable subgroup of G ⁽⁸⁾ [8]. Hence there exists a coordinate system v_1, \dots, v_k of G whose domain includes e and which is adapted to $G(b)^0$, i.e. the slices $[g|v_i(g) = a_i \text{ for all } i = j+1, \dots, k]$ are contained in left cosets of $G(b)^0$ and the slice S^0 with all these $a_i = 0$ is contained in $G(b)^0$. Now consider any $c \in B(b)$ and we shall define some coordinate systems on $B(b)$ at c . Let x_1, \dots, x_d be any coordinate system of M whose domain contains $m' = \pi c$ and $\psi: P \times G \rightarrow B$ any canonical strip map with respect to these taking $(m', e) \rightarrow c$. Then $\psi(P \times S_0) \subseteq B(b)$. The coordinates x_1, \dots, x_d on P and v_1, \dots, v_j (the v_i and j as above) induce coordinates in the usual way on $P \times S_0$ which are carried over to coordinates on $\psi(P \times S_0)$ by ψ . We should not yet refer to these functions on $\psi(P \times S_0)$ as coordinates but it can be shown in routine fashion that such systems of functions on $B(b)$ (depending on choices of $x_1, \dots, x_d, v_1, \dots, v_k, c, \psi$) satisfy the usual compatibility conditions and cover $B(b)$ so they give rise to a unique differentiable structure in which they are coordinate systems. Obviously $B(b)$ is a submanifold of B .

LEMMA 3. If $Q \in \mathcal{Q}, c \in B(b), d \in B(b), Q(c) \in B(b)_c$, then $Q_d \in B(b)_d$.

Proof. We begin with two special cases. The first is where c and d lie on the same fiber, so $d = cg$ and because c and d are in $B(b)$ we have $g \in G(b)$. Consider any strip map $\phi: O \times G \rightarrow B$ such that $\phi(m, e) = c$. It is trivial that $\phi \circ L_g \circ \phi^{-1}$ maps $B(b) \cap \pi^{-1}(0)$ diffeomorphically onto itself and takes $c \rightarrow d, Q(c) \rightarrow Q(d)$. Hence $Q(d) \in B(b)_d$ in this case. The second special case is where there exists a coordinate system x_1, \dots, x_d of M with center at $m = \pi c$ and a canonical (with respect to these x_i) strip map $\psi: P \times G \rightarrow B$, such that $d = \psi(m', e)$ for some $m' \in P$. Let v_1, \dots, v_k be coordinates of G on a neighborhood of e , adapted to $G(b)^0$, so we get coordinates $y_1, \dots, y_d, y_{d+1}, \dots, y_{d+j}$ (these are the coordinates obtained from x_1, \dots, x_d and v_1, \dots, v_k as in the definition of the differentiable structure on $B(b)$) of $B(b)$ whose domain is a $B(b)$ -neighborhood of c . Because $Q \in \mathcal{Q}$ and $Q(c) \in B(b)_c$ we have $Q(c) = \sum_{i=d+1}^{d+j} a_i Y^i(c)$, hence $Q(d) = \sum_{i=d+1}^{d+j} a_i Y^i(d)$, showing $Q(d) \in B(b)_d$. The

⁽⁷⁾ L_h denotes left multiplication by h on G .

⁽⁸⁾ That $G(b)^0$ is a Lie group was pointed out to us by Chevalley, who also found Yamabe's theorem independently.

lemma now follows in general for any $c, d \in B(b)$ by a sequence of steps involving these special cases.

LEMMA 4. $G_0(b) = G(b)^0$.

Proof.⁽⁹⁾ Clearly $G_0(b) \subseteq G(b)^0$ and since both are arcwise connected both are differentiable subgroups of G ; hence $G_0(b)$ is a differentiable subgroup of $G(b)^0$. Now we show the existence of a mapping of $\pi_1(M)$ onto the left cosets of $G_0(b)$ in $G(b)^0$. Since $\pi_1(M)$ contains only countably many elements this will prove the left cosets of $G_0(b)$ in $G(b)^0$ are at most countable in number. This will imply $G_0(b)$ is open in $G(b)^0$ and hence, by connectedness, that $G_0(b) = G(b)^0$.

For each $\alpha \in \pi_1(M)$ let $E(\alpha) = [g \in G(b)^0 \mid \text{there exists a piecewise differentiable horizontal curve from } b \text{ to } bg \text{ whose projection into } M \text{ is in the homotopy class } \alpha]$. Each $E(\alpha)$ is contained in a left coset of $G_0(b)$, for if g_1 and g_2 are in $E(\alpha)$ let ρ_i be a piecewise differentiable horizontal curve from b to bg_i ; then $(\rho^{-1} \circ R_{\rho_2^{-1}g_1}) \cdot \rho_1$ is a piecewise differentiable horizontal curve from b to $bg_2^{-1}g_1$ and its projection into M is $\pi(\rho_2^{-1} \cdot \rho_1)$, showing $g_2^{-1}g_1 \in G_0(b)$. Since each $g \in G(b)^0$ is in some $E(\alpha)$ this furnishes the desired mapping of $\pi_1(M)$ onto left cosets.

If H is a connected differentiable subgroup of a Lie group G with only countably many left cosets, then H is open in G because otherwise H would be of measure 0 (H is trivially a measurable subset of G with respect to the Haar measure of G because, choosing coordinates adapted to H on a neighborhood of e in G , we see some H -neighborhood of e is measurable, and then H is a countable union of translates of this neighborhood) so G itself would have measure 0.

Proof of Theorem 2. We first describe, for each $c \in B(b)_e$, a subspace Δ_c of B_c , then prove $\Delta_c = B(b)_e$, from which the theorem will follow. Note that, trivially, $H_c \subseteq B(b)_e$ for $c \in B(b)$.

Consider any horizontal vector fields K and K' defined on any open submanifold O of B and for each $d \in O \cap B(b)$ consider the vector field $Q \in \mathcal{Q}$ generated by $V([K, K'](d))$ —so Q is defined on all of B and depends on K, K' and d . We do this for all such K, K', d , getting a subset \mathcal{R} of \mathcal{Q} , and define $\mathcal{L}(b)$ to be the Lie algebra of vector fields on B generated by \mathcal{R} . Clearly $\mathcal{L}(b) \subseteq \mathcal{Q}$, and we shall prove later that $\mathcal{L}(b) = q(L(b))$. We define Δ_c , for $c \in B(b)$, to be the subspace of B_c spanned by H_c and all Q_c for which $Q \in \mathcal{L}(b)$.

Now we prove $\Delta_c \subseteq B(b)_e$ for all $c \in B(b)$. Clearly if K is any horizontal vector field on an open subset of B , then $K(c) \in B(b)_e$. Hence if K_1 and K_2 are two such and K'_1, K'_2 are their restrictions to $B(b)$, then $[K'_1, K'_2](c) = [K_1, K_2](c)$, so $B(b)_e$ contains $[K_1, K_2](c)$, and since it contains all horizontal vectors at c , $B(b)_e$ contains $V([K_1, K_2](c))$. This with Lemma 3 shows

⁽⁹⁾ This lemma is an extension of one due to Borel and Lichnerowicz [1], and the proof is a variation on theirs.

$B(b)_c$ contains also $Q(c)$ for any $Q \in \mathcal{Q}$ and $d \in B(b)$ with $V([K_1, K_2](d)) = Q(d)$. Finally, if $Q_1, Q_2 \in \mathcal{Q}$ and $Q_1(c), Q_2(c) \in B(b)_c$ then $[Q_1, Q_2](c) \in B(b)_c$, for if Q'_1, Q'_2 are their restrictions to $B(b)$ then Lemma 3 shows $Q'_1(d), Q'_2(d) \in B(b)_d$ for all $d \in B(b)$, hence $[Q_1, Q_2](c) = [Q'_1, Q'_2](c) \in B(b)_c$. This proves $\Delta_c \subseteq B(b)_c$ for all $c \in B(b)$.

Thus Δ is a distribution on the manifold $B(b)$. We prove now that Δ is involutive and has a differentiable base in the neighborhood of any point of $B(b)$. Let Q_1, \dots, Q_r be any base for $\mathcal{L}(b)$ and K_1, \dots, K_d any differentiable base for H in the neighborhood of any $c \in B(b)$. Let Q'_i and K'_i denote the restrictions of Q_i and K_i to $B(b)$. The Q'_i and K'_i are clearly a differentiable base for Δ throughout a neighborhood of c and to show Δ involutive it is sufficient, by Proposition 1, p. 88 of [6] to show the bracket of any pair of elements from such a base belongs to Δ . But $[Q'_i, Q'_j]$ belongs to Δ because $\mathcal{L}(b)$ is closed under brackets, $[K'_i, K'_j]$ belongs to Δ from the definition of Δ_d , and $[Q'_i, K'_j]$ belongs to Δ because Lemma 1 of §1 implies it is horizontal.

Now we show $\Delta_c \supseteq B(b)_c$, hence $\Delta_c = B(b)_c$, for all $c \in B(b)$. Let ρ be a piecewise differentiable horizontal curve in $B(b)$ from b to c . Because it is horizontal, Δ is involutive, and $\Delta_d \supseteq H_d$, we can prove ρ lies in the integral manifold D of Δ through b . This can be done by extending the tangent vectors to ρ to a differentiable horizontal vector field K on a neighborhood of b ; then K , when restricted to D , must have an integral curve through b in B and by uniqueness of integral curves in $B(b)$ this must be ρ . This shows $D \supseteq B(b)$, hence $\Delta_c \supseteq B(b)_c$.

Now note, for any $c \in B(b)$, that if s and t are in H_c and K_1, K_2 are any horizontal differentiable vector fields on a neighborhood of c with $K_1(c) = s, K_2(c) = t$, then $\Omega(s, t) = -\omega([K_1, K_2](c))$. This holds because the formula for $\Omega = D\omega$ as the "horizontal part" of the exterior derivative gives, for any vector fields W_1, W_2 on B and linear functional f on L ,

$$f(D\omega(W_1(c), W_2(c))) = (HW_1(c))(f \circ \omega)(HW_2) - (HW_2(c))(f \circ \omega)(HW_1) - (f \circ \omega)([HW_1, HW_2](c)).$$

Because ω is vertical, i.e. vanishes on all horizontal vectors, this reduces to

$$f(D\omega(W_1(c), W_2(c))) = - (f \circ \omega)([HW_1, HW_2](c))$$

and hence

$$D\omega(W_1(c), W_2(c)) = - \omega([HW_1, HW_2](c)).$$

This trivially implies the above statement. Hence, for any fixed $c \in B$, the set of all such $\Omega(s, t)$ is the set of all $\omega(V[K_1, K_2](c))$. Thus the set R of all $\Omega(s, t)$ where s and t run through $B(b)_c$ and c runs over $B(b)$ is the set of all $\omega(V[K_1, K_2](c))$ where K_1 and K_2 run through all horizontal vector fields defined on a B -neighborhood of c and c runs over $B(b)$. The set of all these $V[K_1, K_2](c)$ is the same as the set of all $Q(c)$ where $Q \in \mathcal{R}$ and $c \in B(b)$ and,

since $\omega(Q(c)) = \omega(Q(d))$, Lemma 3 implies \mathcal{R} is the set of all $\omega(Q(b))$ (b fixed). Thus $q(\mathcal{R}) = \mathcal{R}$ and hence $q(L(b)) = \mathcal{L}(b)$. Our definition of the differentiable structure on $B(b)$ shows, for ϕ any strip map taking $(m, e) \rightarrow b$ ($m = \pi b$), that ϕ_m takes the set of all $A(e)$, where A is in the Lie algebra of $G(b)^0$, onto the vertical part of $B(b)$. Hence, by what was proved above, ϕ_m takes the set of all such $A(e)$ onto the vertical part of Δ_b , i.e. onto the set of all $Q(b)$ where $Q \in \mathcal{L}(b)$. Because $q(L(b)) = \mathcal{L}(b)$ this shows that $L(b)$ is the Lie algebra of $G(b)^0$. Theorem 2 now follows from Lemma 4.

COROLLARY. *If L' is the subalgebra of L generated by all $\Omega(s, t)$ where s and t run through all pairs of tangent vectors at all points of B , then the subgroup of G generated by L' is the component of the identity of the normal subgroup of G generated by all the $G(b)$ for $b \in B$.*

BIBLIOGRAPHY

1. A. Borel and A. Lichnerowicz, *Groupes d'holonomie des variétés riemanniennes*, C.R. Acad. Sci. Paris vol. 234 (1952) pp. 1835-1837.
2. E. Cartan, *Les groupes d'holonomie des espaces généralisés*, Acta Math. vol. 48, pp. 1-42.
3. H. Cartan, *Notions d'algèbre différentielle; application aux groupes de Lie et aux variétés où opère un groupe de Lie*, Colloque de topologie, Bruxelles, 1950, pp. 15-27.
4. ———, *La transgression dans un groupe de Lie et dans un espace fibre principal*, Colloque de topologie, Bruxelles, 1950, pp. 57-71.
5. S. S. Chern, *Notes on differential geometry*, Institute for Advanced Study notes, 1951.
6. C. Chevalley, *Lie groups*, Princeton University Press, 1946.
7. C. Ehresmann, *Les connexions infinitésimales dans un espace fibre différentiable*, Colloque de topologie, Bruxelles, 1950, pp. 29-55.
8. H. Yamabe, *On an arcwise connected subgroup of a Lie group*, Osaka Math. J. vol. 2 (1950) pp. 13, 14.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY,
CAMBRIDGE, MASS.